

**Derogation for Fenamiphos  
2016/17**

[Derogation compiled by the Deciduous Industry Crop Protection Advisory Group (CPAG)]

<b>Crop</b>	<b>Peaches and Nectarines only.</b>		
<b>Country of Origin</b>	South Africa		
<b>Supplier</b>			
<b>Chemical (ai)</b>	Fenamiphos	<b>Trade name/s</b>	Nemacur 400 EC Fenamiphos 400 EC Bonfire 400 EC Atador
<b>Pest/Disease</b>	Nematodes		
<b>Rationale for use</b>	For the control of nematodes in the soil. Sometimes applied in orchards, as a soil treatment but only if necessary according to soil analyses by Nematode lab.		
<b>Alternative, either cultural or chemically?</b>	<b>Chemical</b> : Cadusfos and Furfural These chemicals must be alternated with Fenamiphos for resistance management. Research in progress on non chemical treatment.		
<b>What prevention methods are available?</b>	None		
<b>Dose rate requested</b>	2.5 ml/M <sup>2</sup>	<b>% of full rate?</b>	- 100%
<b>No of applications requested?</b>	Non bearing trees – 2. Bearing trees - 1	<b>Timing i.e. interval between applications</b>	<b>Non bearing trees</b> within 6 weeks after establishment and repeated 6 months later. <b>Bearing trees</b> post harvest.
<b>Harvest Interval</b>	100 days.	<b>Stewardship and engineering mechanism that prevent environmental issues</b>	Precautions and warnings as per label.
<b>Time scale for exit</b>	As soon as a suitable alternative becomes available.		